

there are acupuncture meridians and autonomic nerve fibers in the mouth that are better than injection sites anywhere else in the body.

The point is however that Oral Administration is not a decrease in efficacy of the remedies and the FDA may in fact without realizing it be pushing us toward a more desirable therapeutic goal.

How than do we manage our therapy???

At a recent HEEL seminar in May 2000 at Los Angeles California, the head of HEELs Scientific department from Germany, Dr. Med. Kusterman gave a Lecture on a technique for allergies known as THE IMHAUSER technique that has relevance to this subject mainly because it is effective for allergy but secondarily because REVERSE IMHAUSER (as developed by your author) is the oral form of AUTO-SANGUIS therapy "5 shots".

The general theory behind injectable 5 shots and the REVERSE SHELTON-IMHAUSER technique (which will be described momentarily) is that our immune systems are depressed and need to be stimulated. This therapy of progressive dilutions of blood progressively ramps up the immune response.

Dr. Kusterman postulated that allergy represents a hyper-reactive immune state and needs to be slowed or ramped DOWN (Down regulated). The Imhauser technique (named after Dr. Kusterman's allergist, Dr. Imhauser) is performed as follows:

To start we need the following supplies:

A. six- 6cc rubber-stoppered sterile glass bottles as used in auto-Sanguis therapy

Six- 6cc dropper bottles

B. a supply of 20% ethanol homeopathic diluent (our office buys Everclear at liquor stores and dilutes it 4:1 with bottled water) (4:1 gives a 20% solution)

C. Allergy syringes to transfer 1/2cc amounts of fluids

D. Stock bottles or ampoules of the following remedies
Galium-Heel

Viscum Album Forte
Ubichinon Compositum
Mucosa Compositum
Co-Enzyme Compositum
Lymphomyosot

Start by filling the 6 rubber stoppered bottles with 5cc of the 20% diluent and label them one through six.

Then fill the dropper bottles as follows:

Each ampoule type has a different amount of solution in it. Add each ampoule to a separate bottle AND THAN add enough of the 20% diluent so that each bottle contains a total of 5cc of solution.

Label the bottle with Galium Heel #7 SIXTH WEEK
Label the bottle with Viscum Forte #8 FIFTH WEEK
Label the bottle with Ubichinon #9 FOURTH WEEK
Label the bottle with Mucosa Comp #10 THIRD WEEK
Label the bottle with Co-Enzyme Comp #11 SECOND WEEK
Label the bottle with Lymphomyosot #12 FIRST WEEK

Draw 0.1cc of the patients blood and add it to bottle #1—succuss it 10 times, than withdraw ½cc from the succussed bottle #1 and add it to:

Bottle #2---succuss it ten times, withdraws 1/2cc and adds it to:

Bottle #3---succuss it ten times, withdraws 1/2cc and adds it to:

Bottle #4---succuss it ten times, withdraws 1/2cc and adds it to:

Bottle #5---succuss it ten times, withdraws 1/2cc and adds it to:

Bottle #6---succuss it ten times, withdraws 1/2cc and add it to:

The first dropper bottle

Bottle#7(the one with Galium)---succuss it ten times, withdraw 1/2cc, add to:

Bottle #8(Viscum)---succuss it ten times, withdraws 1/2cc and adds it to:

Bottle #9(Ubichinon)---succuss it ten times, withdraws 1/2cc and adds it to:

Bottle#10(Mucosa)---succuss it ten times, withdraws 1/2cc and adds it to:

Bottle #11(Co-enzyme)---succuss ten times, withdraw 1/2cc, add to:

Bottle #12(Lymphomyosot)---succuss ten times----END OF PROCESS